

How to use

Nyxoïd® Naloxone Nasal Spray

as part of an opioid overdose response



Step 1

Safety

Check the person and area for anything unsafe.



Step 2

Opioid overdose? Look for the signs

- Not responding
- Pinned pupils
- Not breathing normally - shallow slow breathing or snoring
- Pale or cold skin, blue or grey lips and fingertips



Step 3

Call Triple Zero (000)
and put on speaker phone while you
administer Nyxoïd®



"I think it's an
opioid overdose"

Step 4

**Lie person on back and check their nose
is clear**



adis

**24/7
ALCOHOL
AND DRUG
SUPPORT**

adis.health.qld.gov.au

1800 177 833
(free call)




Step 5

Hold Nyxoid® like this

Remember Nyxoid® contains 1 dose only.


Do not press to test!



Step 6

Give Naloxone into nose

Insert Nyxoid® nozzle in nostril. Press until it clicks to give the dose.



Step 7

Are they breathing?

If not, use rescue breathing and/or chest compressions.

(2 breaths / 30 compressions)




Step 8

Repeat dose if needed

If no response, give 2nd dose of Nyxoid® in other nostril.


If available, give another dose every 2 minutes until the person recovers or ambulance arrives.



Step 9

Recovery

Put person in 'recovery position' if you know how. Stay with them until ambulance arrives.



Nyxoid® takes 2-5 minutes to start working and lasts 30-90 minutes.

Need more information?

For information and support call Adis 24/7 Alcohol and Drug Support on **1800 177 833** or scan the QR code to visit our website.

